Selected Topics in Algorithmics, SS15 Exercise Sheet "1": Quicksort and Trapezoidal Decomposition University of Bonn, Department of Computer Science I

- Written solutions have to be prepared until Wednesday 29th of April, 14:30 pm. There is a letterbox in front of room E.01 in the LBH building.
- You may work in groups of at most two participants.

Exercise 1: Probability Space

(4 Points)

Consider a standard 52-card deck of poker cards. Assume we do not distinguish cards by their suits, i.e., cards with the same number are identical. We want to select 5 cards from the deck. Please define the probability space as follows.

- 1. Please describe the sample space Ω . (The outcome can be classified into 6 category, and each category has different number of elements.)
- 2. Please describe the family \mathcal{F} of events, e.g., the total number.
- 3. Please describe the probability function Pr by illustrating the probability for the elements in the sample space. (just one element for each category)
- 4. Let X be the random variable representing the sum of 5 cards. Please compute the expectation of X.

Exercise 2: Average Complexity of Sorting (4 Points)

Given a set N of n real numbers, please analyze the average complexity for the following sorting algorithms over all the n! permutation sequences of N.

- Insertion Sort
- Merge Sort
- Quick Sort (always select the first element)

Exercise 3: Vertical Trapezoidal Decomposition (4 Points)

Given a set N of n line segments with a total number k of intersection in the plane, let S_1, S_2, \ldots, S_n be a random sequence of N, and let N^i be $\{S_1, S_2, \ldots, S_i\}$. Please prove the following.

- 1. The vertical trapezoidal decomposition H(N) of N has O(n+k) trapezoids (faces) even if more than two line segments can intersect at the same point.
- 2. The expected number of trapezoids in $H(N^i)$ is $O(i + ki^2/n^2)$. (Hint: the expected number of intersections)