

Discrete and Computational Geometry, WS1516
Exercise Sheet “6”: Order- k Diagrams University of
Bonn, Department of Computer Science I

- *Written solutions have to be prepared until **Wednesday 16th of December, 12:00 pm**.*
- *There is a letterbox in front of Room E.01 in the LBH building.*
- *You may work in groups of at most two participants.*

Exercise 13: Numbers of vertices, edges, and faces of $V_k(S)$ (8 points)

Let S be a set of n point sites in the Euclidean plane satisfying a general position assumption that no three sites are on the same line and no four sites are on the same circle. For $1 \leq i \leq n - 1$, let $N_i, E_i, I_i, \mathcal{S}_i$ be the numbers of faces, edges, vertices, and unbounded faces of $V_i(S)$, respectively, and let \mathcal{S}_0 be 0. Please prove the following:

1. $E_k = 3(N_k - 1) - \mathcal{S}_k$ and $I_k = 2(N_k - 1) - \mathcal{S}_k$. (Hint: Euler formula. Due the general position assumption, the degree of a Voronoi vertex is 3). (2 points)
2. $N_1 = n$, and $N_2 = 3(n - 1) - \mathcal{S}_1$, and $N_k = 3(N_{k-1} - 1) - \mathcal{S}_{k-1} - 2 \sum_{i=1}^{k-2} (-1)^{k-2-i} (2(N_i - 1) - \mathcal{S}_i)$ implies

$$N_k = 2k(n - k) + k^2 - n + 1 - \sum_{i=0}^{k-1} \mathcal{S}_i.$$

(Hint: By induction on k) (6 points)

Exercise 14: Relation between $V_i(S)$ and $V_{i+1}(S)$ (4 points)

Assume $\text{VR}_i(H, S)$ has m adjacent regions $\text{VR}_i(H_j, S)$, $1 \leq j \leq m$. Let Q be $\bigcup_{1 \leq j \leq m} H_j \setminus H$. Prove that $V_{i+1}(S) \cap \text{VR}_i(H, S) = V_1(Q) \cap \text{VR}_i(H, S)$. (Hint: prove that for all site $r \in (S \setminus H) \setminus Q$, $\text{VR}_1(r, S \setminus H) \cap \text{VR}_k(H, S) = \emptyset$. You can first assume the contrary that $\exists r \in (S \setminus H) \setminus Q$, $\text{VR}_1(r, S \setminus H) \cap \text{VR}_k(H, S) \neq \emptyset$, and then show that it will lead to a contradiction. For any point $x \in \text{VR}_1(r, S \setminus H) \cap \text{VR}_k(H, S)$, \overline{rx} will intersect a Voronoi edge e between $\text{VR}_i(H, S)$ and $\text{VR}_i(H_j, S)$ for some $j \in \{1, \dots, m\}$. Let y be the intersection point between \overline{rx} and e . Discuss nearest neighbors of y , which will lead to a contradiction from the viewpoint of e and the viewpoint of $\text{VR}_1(r, S \setminus H)$.)

Bonus 3: Voronoi edges of k^{th} -order Voronoi diagrams (4 points)

Consider a Voronoi edge e between two adjacent Voronoi regions $\text{VR}_k(H_1, S)$ and $\text{VR}_k(H_2, S)$, where S is a set of n point sites in the Euclidean plane. Please prove the following.

1. $|H_1 \setminus H_2| = |H_2 \setminus H_1| = 1$
2. The circle centered at a point x in e and touching p and q , where $H_1 \setminus H_2 = \{p\}$ and $H_2 \setminus H_1 = \{q\}$, encloses exactly $k - 1$ sites of S .

(Hint: Consider $\text{VR}_{k-1}(H, S)$ and $V_1(S \setminus H)$, where $e \cap \text{VR}_{k-1}(H, S) \neq \emptyset$.)